

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
31 December 2003 (31.12.2003)

PCT

(10) International Publication Number
WO 2004/001120 A1

(51) International Patent Classification⁷: D06L 1/00

(21) International Application Number:
PCT/GB2002/002846

(22) International Filing Date: 24 June 2002 (24.06.2002)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): IMPE-
RIAL CHEMICAL INDUSTRIES PLC [GB/GB]; 20
Manchester Square, London W1U 3AN (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MOTSON, Harold,
Russell [GB/GB]; 10 Grantham Road, Norton, Stock-
ton-on-Tees, Cleveland TS10 1PP (GB). IRVINE,
Derek, John [GB/GB]; 24 Burnmoor Drive, Eaglescliffe,
Stockton-on-Tees TS16 0HZ (GB). APPLEMAN, Eric
[NL/NL]; Adriaan van Bleijenurgstraat 22, NL-3311 LB
Dordrecht (NL).

(74) Agents: ROBERTS, Jonathan, Winstanley et al.;
Uniqema Intellectual Property Dept., Wilton Centre,
Wilton, Redcar TS10 4RF (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



WO 2004/001120 A1

(54) Title: METHOD FOR CLEANING TEXTILES

(57) Abstract: Detergent or micelle free cleaning media based on detergent free and/or micelle free liquid CO₂ and including from 0.01 to 5 % by weight of the formulation of a cleaning additive which is at least one multi-ester having a molecular weight of not more than 750 can be used in dry cleaning of textiles. Desirable cleaning additives are of the formula (1): R¹ (XR²)_n, where X, R¹, R², and n have defined meanings, particularly to be esters of multi-carboxylic acids and mono-hydroxy alcohols or esters of mono-carboxylic acids and multi-hydroxy alcohols.

METHOD FOR CLEANING TEXTILES

This invention relates to cleaning textile materials and products including clothes using liquid carbon dioxide (CO₂) and cleaning additives.

5 The dry cleaning of clothes using fluid carbon dioxide, either as liquid or supercritical fluid, is known from many patents. An early suggestion is in US 4012194 (Maffei) which teaches simply using liquid carbon dioxide as a substitute for halocarbon solvents e.g. perchlorethylene (perc), used in conventional dry cleaning. Later patents develop approaches using detergent materials, including US Patents US 5676705, US 5683473, US 5683977, US 6131421, US 6148644, and US 6299652 assigned to Unilever and their equivalents, which relate to the use of defined detergents based on
10 various classes of polymers and a series of cases, including US Patents 5858022, 6200352, 6280481, 6297206, 6269507 and US published application 200106053 A, assigned to MiCell and their equivalents. Also US 5279615 assigned to Chlorox Co uses cleaning non-polar organic cleaning adjuncts, especially alkanes, in densified, particularly supercritical CO₂.

15 This invention is based on a liquid CO₂ dry cleaning medium including relatively polar multi-esters as cleaning additives which improve the cleaning performance of the liquid CO₂ and give improved handling characteristics as compared with the use of detergents available for use with liquid CO₂. The multi-esters are compounds having 2 or more carboxylic acid ester groups, and molecular weights of not more than 750.

20 The invention accordingly provides a detergent free dry cleaning medium based on liquid CO₂ and including from 0.01 to 5% by weight of the cleaning medium of a cleaning additive which is at least one multi-ester having a molecular weight of not more than 750.

Alternatively the invention provides a micelle free dry cleaning medium based on liquid CO₂ and including from 0.01 to 5% by weight of the cleaning medium of a cleaning additive which is at least one multi-ester having a molecular weight of not more than 750.

25 The invention includes a method of dry cleaning which includes contacting textile material, particularly clothes, with a detergent free dry cleaning medium based on liquid CO₂ and including from 0.01 to 5% by weight of the cleaning medium of a cleaning additive which is at least one multi-ester having a molecular weight of not more than 750.

30 Further alternatively, the invention includes a method of dry cleaning which includes contacting textile material, particularly clothes, with a micelle free dry cleaning medium based on liquid CO₂ and including from 0.01 to 5% by weight of the cleaning medium of a cleaning additive which is at least one multi-ester having a molecular weight of not more than 750.

In the present invention in describing cleaning media as "detergent free" we mean that they do not include amphiphilic materials that aid soil removal from textiles. In describing cleaning media as
35 "micelle free" we mean that the cleaning medium does not contain micelles of cleaning additives.

We have found that the presence of detergents including those which may form micelles in liquid CO₂ can reduce the effectiveness of the cleaning additives used in the invention.

The cleaning additive multi-esters used in this invention are desirably of the formula(I):



5 where

X is -C(O)O- or -OC(O)- ; such that

where X is -C(O)O-,

R¹ is a direct bond or the residue of a C₁ to C₁₀ hydrocarbyl group from which n hydrogen atoms have been removed; and

10 R² is a C₁ to C₁₀ hydrocarbyl group; and

where X is -OC(O)-,

R¹ is or the residue of a C₂ to C₁₀ hydrocarbyl group from which n hydrogen atoms have been removed; and

R² is H or a C₁ to C₁₀ hydrocarbyl group; and

15 n is from 2 to 5;

the compound having a molecular weight of not more than 750.

These cleaning additive multi-esters can be divided into two sub-classes respectively of the formulae (Ia) and (Ib) below. Compounds of the formula (Ia) are esters of a multi-carboxylic acid and a mono-hydroxy alcohol:



where

X is -C(O)O-;

R^{1a} is a direct bond or the residue of a C₁ to C₁₀ hydrocarbyl group from which n hydrogen atoms have been removed;

25 R^{2a} is a C₁ to C₁₀ hydrocarbyl group; and

the compound having a molecular weight of not more than 750.

Examples of compounds of the formula (Ia) include di-esters of dicarboxylic acids such as succinic, glutaric and adipic acids.

Compounds of the formula (Ib) are esters of a monocarboxylic acid and a multi-hydroxy alcohol:



where

X is -OC(O)- ;

R^{1b} is or the residue of a C₂ to C₁₀ hydrocarbyl group from which n hydrogen atoms have been removed; and

35 R^{2b} is H or a C₁ to C₁₀ hydrocarbyl group; and

the compound having a molecular weight of not more than 750.

Examples of compounds of the formula (Ib) include esters of multi-hydroxyl compounds such as triacetin (glycerol triacetate), ethylene glycol diacetate and pentaerythritol tetra-acetate.

The precise mode of action of the multi-ester cleaning additives is not clear. They do appear to boost the overall cleaning performance of liquid CO₂ but operating at levels that are significantly lower that would be expected to be effective if the effect were simply additive co-solvency. In addition the use of these additives gives improved handling of textiles cleaned using them as compared with no cleaning additives or commercially available detergents for use in liquid CO₂.

Within the formula (I) above, generally is desirable that the group X is -C(O)O- as these compounds seem to provide superior effects in cleaning. Among such compounds, the group R¹ is desirably -(CH₂)_m- where m = 2 to 6, particularly 2 to 4 and especially as in the mixed ester of succinic, glutaric and adipic acids; and the group R² is desirably methyl, ethyl or propyl, particularly methyl. Thus, the dimethyl esters of succinic, glutaric and adipic acids, particularly as mixtures are particularly desirable additives.

The molecular weight of the cleaning additive is not more than 750 and is desirably not more than 500. In practice the molecular weight for individual components e.g. of formula (I) can be as low as 118 (dimethyl oxalate) but will not usually be lower than 146 (dimethyl succinate and ethylene glycol diacetate). More usually on average the molecular weight will be at least 150, particularly from 150 to 300. The mixed dimethyl esters of succinic, glutaric and adipic acids can have molecular weights ranging from about 150 to 170 e.g. for an approximately 1:1:3 mixture the average molecular weight is about 165.

In order to maintain the desired high polarity, the ratio of oxygen to carbon atoms in the molecules of the cleaning additive is (on average) desirably from from 1:1 to 1:5 particularly from 1:1 to 1:3 and especially from 1:1 to 1:1.5. The mixed dimethyl esters mentioned above have an average ratio of ca 1:1.23.

The amount of cleaning additive multi-ester present in the cleaning medium is from 0.01 to 5%, usually from 0.05 to 2%, more usually from 0.1 to 1%, particularly from 0.1 to 0.5% and more particularly from 0.1 to 0.3% by weight of the cleaning medium. The use of lower amounts of cleaning additive will not generally give useful results and use of larger amounts does not appear to give additional benefits and may result in including so much additive in the system that additive residues are deposited onto the textiles being cleaned or left on the walls of the cleaning apparatus.

Other ingredients can be included in the dry cleaning formulation such as fragrances, optical brighteners, fabric conditioners such as softeners, and sizes e.g. starch, enzymes, bleaches, particularly peroxide bleaches e.g. organic and/or inorganic peroxides or hydrogen peroxide or a source of hydrogen peroxide.

The textiles to be cleaned will usually be garments and can be of woven or non-woven fabrics. The fibre making up the fabric can be or include a wide range of natural and synthetic fibres including polyamides particularly natural polyamides such as silk and wool and synthetic polyamides such as nylon, cellulosic fibres such as cotton, linen and rayon, synthetic polymers such as polyester, particularly polyethylene terephthalate or related copolymers, or acetate polymers. When fabrics including acetate polymers and possibly nylon polymers are cleaned it is best to avoid direct contact between the fabric and high concentrations of or neat multi-ester additives. When neat or at high concentration, the multi-ester additives may swell or dissolve such polymers leading to fabric damage. Thus it is desirable to pre-mix the multi-ester with CO₂ before permitting contact with such polymers. Pre-mixing the multi-ester cleaning additive with CO₂ to give a concentration of less than about 10%, more usually less than 5%, and desirably not more than 2% by weight of the cleaning additive in the liquid CO₂ based cleaning medium before the additive comes into contact with the textile seems to avoid this potential problem.

The particular mode of operation will depend on the equipment used. Generally the cleaning will be carried out in a drum, which may have its axis vertical or horizontal. The textiles are introduced into the drum which is then sealed and filled with the cleaning medium including carbon dioxide typically to give a mixture of liquid and gaseous CO₂ in the drum. The textiles and liquid CO₂ based cleaning medium are then agitated to give thorough mixing and contact between the cleaning medium and textiles. The textiles will be contacted with the cleaning medium for a time adequate to clean the textiles to the desired extent. The cleaning medium is then separated from the textiles, typically by draining or venting it from the drum. Generally the textiles will be subject to one such cleaning cycle, but if desired the cleaning cycle may be repeated to obtain a higher degree of cleaning. Usually, the textiles are subject to at least one rinse cycle with liquid carbon dioxide usually not including cleaning additives, but which may include fabric softeners, optical bleaches etc if desired. The rinse liquid is similarly separated from the textiles, which can then be recovered by de-pressurising the drum and opening it to remove the textiles.

Any suitable apparatus for dry cleaning with liquid carbon dioxide can be used. Typically such apparatus includes a drum in which the cleaning is carried out. The drum may have its axis horizontal or vertical. (Other angles of orientation will generally be less convenient in operation.) Providing agitation in a horizontal axis drum can simply be by rotation around its axis. Vertical axis drums will usually include an agitator which can be moved to agitate the drum contents. Other means of agitation include paddles or vanes in the drum or by jetting liquid CO₂ into the mixture of cleaning medium and textiles in the drum. Suitably vigorous agitation may give rise to cavitation in the cleaning medium and this may improve the cleaning performance.

Typically the cleaning temperature will be from -10 to 25°C, more usually from 5 to 25°C, particularly from 10 to 20°C. The operating temperature will not usually be above about 25°C to maintain the cleaning medium a reasonable margin from the critical point of CO₂, as supercritical

CO₂ may extract textile dyes from fabrics. Operating at or near ambient temperature simplifies operation of the process, but using a lower temperature means that the CO₂ is more dense and a more effective cleaning agent. Temperatures in the range 10 to 17°C, particularly 12 to 15°C generally provide a reasonable balance of properties and are thus advantageous.

- 5 During cleaning the cleaning medium must be kept at a pressure which maintains the CO₂ at least partially as a liquid. This will usually be the vapour pressure of the cleaning medium at the temperature of operation because, as is noted above, it is desirable for both liquid and gaseous CO₂ to be present. At the typical operating temperatures noted above, the corresponding pressures are approximately 2.7 to 6.4 MPa, more usually from 4 to 6.4 Mpa, particularly from
- 10 4.5 to 5.7 Mpa and balancing density and temperature 4.5 to 5.5 Mpa, particularly from 4.9 to 5.1 Mpa.

The invention is illustrated by the following Examples. All parts and percentages are by weight unless otherwise indicated.

Materials

- CA1 mixed esters: dimethyl adipate (ca 60%), dimethyl glutamate (ca 20%), and dimethyl succinate (ca 20%)
- CA1a additive CA1 plus a fragrance
- CA2 mixed esters: dimethyl adipate (ca 90%) and dimethyl glutamate (ca 10%)
- CA3 dimethyl adipate
- CA4 triacetin
- CD1 Kreussler - conventional formulated detergent
- CD2 Fabritech 5565 - conventional formulated detergent
- CD3 Conventional detergent (composition not known)

Cleaning testing used standard "Krefeld" stained cloths. The codes for these cloths include a number indicating the fabric type and a letter or letters indicating the soil as follows:

Cloth Type		Soil Type			
10	cotton	C	WFK soil*/lanolin mix	GM	used motor oil
20	polycotton	D	sebum	TE	clay
30	polyester (PET)	LS	Lipstick	PF	pigment/vegetable fat

- * WFK soil - a mixed soil based on kaolinite and containing soot and iron oxide pigments

Cleaning effectiveness - was assessed spectrometrically (using an X-Rite Spectrophotometric Colour Measurement system) by comparison of commercially available standard soiled cloths before and after cleaning with the results given as % stain removal.

Example 1

- Various cleaning additives were tested for efficacy in removing stains from standard stained cloths using the experimental cleaning machine and method set out below.

Test Cleaning Procedure

- An experimental cleaning machine is based on a pressure cylinder ca 50 cm long by 15 cm diameter (external); internal volume ca 6 l as the cleaning vessel. Connections are provided to enable the cylinder to be filled with carbon dioxide and emptied and for holding test cloths in the vessel.

- Soiled fabric samples are held in place inside the pressure cylinder, the desired additive is introduced into the bottom of the cylinder using a syringe and the cylinder sealed. The cylinder is filled initially with gaseous carbon dioxide (to a minimum of 30 bar pressure) and then the desired quantity, usually from 1.5 to 2.0 kg (measured by logging the weight loss of the supply cylinder), of liquid carbon dioxide is introduced. The supply connections are removed and the test cylinder is rotated end-over end for a predetermined time. The cylinder

is then suspended with its axis vertical so that the 'dirty' liquid drains away from the washed fabric samples under gravity. The 'dirty' liquid CO₂ is vented to atmosphere. A rinse stage is normally carried out by repeating the filling process but without using any cleaning additive. The fabric samples are then removed from the machine removed and the stains examined using a computer controlled spectrophotometric colour measurement system.

The cleaning conditions and the results obtained are set out in Table 1 below

Table 1

Ex No	Wash Time (min)	Rinse Time (min)	Final Press. (Bar)	Temp (°C)	Additive		% Soil Removed				
					type	(%w/w)	30C	30D	10LS	10PF	10GM
1.C.1	15	0	50	16	none	-	25	35	27	23	24
1.C.2	15	15	50/51	13/15	CD1	0.2	28	39	29	29	26
1.1	15	15	50	15	CA1	0.2	36	41	38	28	28
1.2	15	15	45	12	CA1	0.2	33	32	30	23	24
1.3	15	15	48	14	CA2	0.2	20	34	29	19	19
1.4	15	15	-	-	CA3	0.2	33	42	30	25	27

Example 2

Further tests were carried out in commercial scale liquid CO₂ dry cleaning equipment using standard Krefeld soiled cloths, pinned to blank textile sheets to provide more realistic behaviour in the cleaning machine. The results are set out in Table 2 below:

Table 2

Ex No	Additive		30C	30D	20MU	10LS	10PF	10TE	10GM
	type	amount							
2.1.C.1	none	-	42	55	31	36	38	18	26
2.1.C.2	CD2	0.2	20	35	21	32	29	14	22
2.1	CA1a	0.2	48	67	39	39	47	24	28
2.2.C.1	none	-	38	64	32	38	41	18	26
2.2.C.2	CD3	0.2	45	63	33	36	40	22	23
2.2	CA1a	0.2	45	69	32	39	45	23	24

The textiles cleaned using additive CA1a had a significantly improved feel as compared with cloths cleaned with liquid CO₂ alone or using the commercial detergent additives.

Claims

- 1 A detergent free dry cleaning medium based on liquid CO₂ and including from 0.01 to 5% by weight of the cleaning medium of a cleaning additive which is at least one multi-ester having a molecular weight of not more than 750.
- 5 2 A dry cleaning formulation as claimed in claim 1 wherein the multi-ester includes at least one compound of the formula (I):
- $$R^1(XR^2)_n \quad (I)$$
- where
 X is -C(O)O- or -OC(O)- ; such that
 10 where X is -C(O)O-,
 R¹ is a direct bond or the residue of a C₁ to C₁₀ hydrocarbyl group from which n hydrogen atoms have been removed; and
 R² is a C₁ to C₁₀ hydrocarbyl group; and
 where X is -OC(O)-,
 15 R¹ is or the residue of C₂ to C₁₀ hydrocarbyl group from which n hydrogen atoms have been removed; and
 R² is H or a C₁ to C₁₀ hydrocarbyl group; and
 n is from 2 to 5;
 the compound having a molecular weight of not more than 750.
- 20 3 A dry cleaning formulation as claimed in claim 2 wherein the multi-ester is of the formula (Ia):
- $$R^{1a}(XR^{2a})_n \quad (Ia)$$
- where
 X is -C(O)O-;
 R^{1a} is a direct bond or the residue of a C₁ to C₁₀ hydrocarbyl group from which n
 25 hydrogen atoms have been removed;
 R^{2a} is a C₁ to C₁₀ hydrocarbyl group; and
- 4 A dry cleaning formulation as claimed in claim 3 wherein the multi-ester is a dimethyl ester of adipic, glutaric or succinic acids or a mixture of such esters.
- 5 A dry cleaning formulation as claimed in any one of claims 1 to 4 wherein the average
 30 molecular weight of the multi-ester(s) is from 150 to 300.
- 6 A dry cleaning formulation as claimed in any one of claims 1 to 5 wherein the average ratio of oxygen atoms to carbon atoms in the multi-ester(s) is from 1:1 to 1:5.
- 7 A dry cleaning formulation as claimed in claim 6 wherein the average ratio of oxygen atoms to carbon atoms in the multi-ester(s) is from 1:1 to 1:1.5.

- 8 A dry cleaning formulation as claimed in any one of claims 1 to 7 wherein the amount of cleaning additive multi-ester present in the cleaning medium is from 0.1 to 0.5% by weight of the cleaning medium.
- 9 A dry cleaning formulation as claimed in any one of claims 1 to 8 which additionally includes
5 at least one fragrance, optical brightener, fabric conditioner, enzyme and/or bleach.
- 10 A method of dry cleaning which includes contacting textile material with a detergent free dry cleaning medium based on liquid CO₂ and including from 0.01 to 5% by weight of the cleaning medium of a cleaning additive which is at least one multi-ester having a molecular weight of not more than 750.
- 10 11 A method as claimed in claim 10 wherein the multi-ester includes at least one compound of the formula (I): $R^1(XR^2)_n$ where X, R¹, R² and n are as defined in claim 2, the compound having a molecular weight of not more than 750.
- 12 A method as claimed in claim 11 wherein the multi-ester is of the formula (Ia):

$$R^{1a}(XR^{2a})_n \quad (Ia)$$

 15 where
 X is -C(O)O-;
 R^{1a} is a direct bond or a C₁ to C₁₀ hydrocarbyl group from which n hydrogen atoms have been removed;
 R^{2a} is a C₁ to C₁₀ hydrocarbyl group; and
- 20 13 A method as claimed in any claim 12 wherein the multi-ester is a dimethyl ester of adipic, glutaric or succinic acids or a mixture of such esters.
- 14 A method as claimed in any one of claims 10 to 13 wherein the average molecular weight of the multi-ester(s) is from 150 to 300.
- 15 A method as claimed in any one of claims 10 to 14 wherein the average ratio of oxygen
25 atoms to carbon atoms in the multi-ester(s) is from 1:1 to 1:1.5.
- 16 A method as claimed in any one of claims 10 to 15 wherein the amount of cleaning additive multi-ester present in the cleaning medium is from 0.1 to 0.5% by weight of the cleaning medium.
- 17 A method as claimed in any one of claims 10 to 16 which additionally includes at least one
30 fragrance, optical brightener, fabric conditioner, enzyme and/or bleach.
- 18 A method as claimed in any one of claims 10 to 17 wherein the multi-ester is pre-mixed with liquid CO₂ before contacting the textiles.
- 19 A method as claimed in any one of claims 10 to 18 wherein the cleaning process is carried out at a temperature of from -5 to 25°C.

- 20 A method as claimed in claim 19 wherein the temperature is from 5 to 20°C.
- 21 A method as claimed in claim 20 wherein the temperature is from 12 to 15°C.

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/02/02846

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 D06L1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 D06L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 2002/010965 A1 (RACETTE TIMOTHY L ET AL) 31 January 2002 (2002-01-31) page 6, paragraph 154 - paragraph 161 page 10, paragraph 236; claims 1,10,17,18,27,34; tables 1,2 ---	1-17
A	US 5 858 022 A (CAUBLE DAVID F ET AL) 12 January 1999 (1999-01-12) cited in the application column 3, line 17 - line 18 ---	1,10
A	US 6 369 014 B1 (BINDER DAVID ALAN ET AL) 9 April 2002 (2002-04-09) the whole document --- -/--	1,10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

27 May 2003

Date of mailing of the international search report

13/06/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Blas, V

INTERNATIONAL SEARCH REPORT

Internati Application No
PCT/02/02846

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with Indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 00 42249 A (MICELL TECHNOLOGIES INC ;CAUBLE DAVID F (US); MCCLAIN JAMES B (US)) 20 July 2000 (2000-07-20) page 5, line 18 - line 21 page 6, line 20 - line 22 page 7, line 30; claims -----</p>	1,10

INTERNATIONAL SEARCH REPORT

In m on patent family members

Internati Application No

PCT/G 2/02846

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2002010965	A1	31-01-2002	US 6355072 B1 12-03-2002
			WO 02086222 A1 31-10-2002
			AU 8021700 A 30-04-2001
			AU 8021800 A 30-04-2001
			BR 0014770 A 17-09-2002
			CA 2388500 A1 26-04-2001
			CA 2388913 A1 26-04-2001
			EP 1224351 A1 24-07-2002
			EP 1224352 A1 24-07-2002
			JP 2003512111 T 02-04-2003
			NO 20021764 A 14-06-2002
			NO 20021765 A 17-06-2002
			WO 0129305 A1 26-04-2001
			WO 0129306 A1 26-04-2001
			US 2002100124 A1 01-08-2002
			US 2002011258 A1 31-01-2002
US 5858022	A	12-01-1999	AU 736088 B2 26-07-2001
			AU 8922098 A 16-03-1999
			CA 2301636 A1 04-03-1999
			EP 1007779 A1 14-06-2000
			JP 2001514337 T 11-09-2001
			WO 9910585 A1 04-03-1999
			US 6218353 B1 17-04-2001
			US 6200352 B1 13-03-2001
			US 6258766 B1 10-07-2001
US 6369014	B1	09-04-2002	CA 2383480 A1 24-11-2002
			EP 1260574 A2 27-11-2002
			US 6475968 B1 05-11-2002
WO 0042249	A	20-07-2000	US 6200352 B1 13-03-2001
			AU 2732200 A 01-08-2000
			CA 2359837 A1 20-07-2000
			EP 1144748 A1 17-10-2001
			WO 0042249 A1 20-07-2000
			US 6258766 B1 10-07-2001